

Baby Immunisation



Today your child was immunised against disease.

Most immunisations have no reaction, but ..

Too Hot



Baby is too **hot**.



Give them a drink (water or breastfeed)

Loosen their clothes




Crying

Baby is unsettled, grumpy, **crying** or has a fever



Give them paracetamol as advised by the nurse

Give them plenty of hugs



Red or Swollen

Red or **swollen** at the injection site



Place a cool wet cloth on site

Do not rub the site.

Sleepy or Not Feeding



Baby is **sleepy** or slow to **feed**

Watch them carefully



Encourage them to drink




Worried























Worried about your child?



Call your nurse doctor or hospital



Vaccine and Common Side Effects

<input type="checkbox"/>	Diphtheria-tetanus-acellular pertussis (DTaP) - inactive Polio (IPV)				6hrs	
<input type="checkbox"/>	Haemophilus influenzae type b Hepatitis B (Hib-HepB)				6hrs	
<input type="checkbox"/>	Hepatitis B (Hep B)				6hrs	
<input type="checkbox"/>	Measles-mumps-rubella (MMR)					Rash 5-12 days
<input type="checkbox"/>	Haemophilus Influenzae type b (Hib)					6hrs
<input type="checkbox"/>	Meningococcal B (MenZB™)				6hrs	 

Treat the side effects as shown in this leaflet or if you are worried
phone your doctor or nurse, tell them the vaccine and the symptoms.

**Mangere Refugee
Resettlement
Centre
Medical Clinic**

**After a child's
Immunisation**

